



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

# Multiple signalling modalities mediated by dendritic exocytosis of oxytocin and vasopressin

### Citation for published version:

Ludwig, M & Stern, J 2015, 'Multiple signalling modalities mediated by dendritic exocytosis of oxytocin and vasopressin', *Philosophical Transactions of the Royal Society B: Biological Sciences*, vol. 370, no. 1672.  
<https://doi.org/10.1098/rstb.2014.0182>

### Digital Object Identifier (DOI):

[10.1098/rstb.2014.0182](https://doi.org/10.1098/rstb.2014.0182)

### Link:

[Link to publication record in Edinburgh Research Explorer](#)

### Document Version:

Peer reviewed version

### Published In:

Philosophical Transactions of the Royal Society B: Biological Sciences

### General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# Multiple signalling modalities mediated by dendritic exocytosis of oxytocin and vasopressin

Mike Ludwig<sup>1</sup> and Javier Stern<sup>2</sup>

<sup>1</sup>Centre for Integrative Physiology, University of Edinburgh, Edinburgh, UK, <sup>2</sup>Department of Physiology, Medical College of Georgia, Georgia Regents University, Augusta GA, USA

**Keywords:** hypothalamus, supraoptic nucleus, paraventricular nucleus, priming, calcium, homeostasis

## Summary

The mammalian hypothalamic magnocellular neurons of the supraoptic and paraventricular nuclei are among the best understood of all peptidergic neurons. Through to their anatomical features, vasopressin- and oxytocin-containing neurones have revealed many important aspects of dendritic functions. Here we review our understanding of the mechanisms of somato-dendritic peptide release, and the effects of autocrine, paracrine and hormone-like signalling on neuronal networks and behaviour.

## Of secret messages and public announcements.

Forms of information processing and intercellular communication in the brain may be classified, at least in part, according to distinct spatio-temporal features. At one end of the spectrum is classical chemical synaptic transmission. Chemical synapses are structurally organized units with a well-defined physical substrate, and have evolved to transfer information between pairs of neurons efficiently, in a precise, spatially constrained and rapid manner. The strength and time course of this “hard-wired” communication is dependent on the probability of presynaptic transmitter release, the affinity of the postsynaptic receptors for the transmitter, the density of postsynaptic receptors clustered at highly specialized sites, and the rate of diffusion/uptake of the neurotransmitter at/from the synaptic cleft [1-4].

At the opposite end of the spatio-temporal spectrum, paracrine or hormone-like signalling modalities mediate transfer of information between entire populations of neurons, which in some cases may be located relatively distant from each other, acting in a more diffuse, less spatially constrained manner and on a slower time scale. In the hard-wired chemical synapse, the “secrecy” of the communication is largely determined by the spatially constrained structure of the synapse. Conversely, in paracrine transmission, specificity is solely determined by the specificity of the signal/receptor interaction. Examples of signalling mechanisms acting at more distant sites include release of catecholamines and acetylcholine from en passant boutons on axonal segments [5], and gaseous neurotransmitters including nitric oxide and carbon monoxide [6]. However, the prototypes for hormone-like signalling within the brain are many neuropeptides, including vasopressin and oxytocin, released from their somata and dendrites. They are public announcements; they are messages not from one cell to another, but rather a message that is directed from one population of neurons to another [7-9].

## The hypothalamo-neurohypophysial axis – a model system to study dendritic

\*Author for correspondence (mike.ludwi@ed.ac.uk).

†Present address: Centre for Integrative Physiology, University of Edinburgh, George Square, Edinburgh EH8 9XD, UK

## peptide release

The dendrites of magnocellular neurons (MCNs) of the hypothalamic supraoptic nucleus (SON) and paraventricular nucleus (PVN) have some unique characteristics compared to other neurons in the central nervous system. They are aspiny, branch sparsely, in many cases are aggregated in bundles, and are relatively thick and varicose. Dendrites in MCNs are structurally dynamic, undergoing activity-dependent remodelling, including shrinkage/elongation, altered branching patterns and increased bundling [10, 11]. Another salient feature is that in more than 60% of MCNs, axons arise from a dendrite rather than more conventionally from the soma [10, 12]. These axon-bearing dendrites may not only be privileged in their ability to influence spiking initiation and overall neuronal output [13], but they could be in turn more efficiently affected by back-propagating action potentials (see below).

The MCNs of the SON and PVN themselves are large and can easily be identified. Their cell bodies and dendrites are aggregated into compact and homogenous nuclei located in and receiving input from the central nervous system. Their axons project to the posterior pituitary gland, which lacks an effective blood-brain barrier allowing secretion from this site to enter the systemic circulation. MCNs dendrites are known to store the majority of the neuropeptide content present in the SON and PVN, and studies of dendritic release using push-pull perfusion or microdialysis [14, 15] can be accomplished without contamination by local synaptic release or reuptake of peripherally-released peptides (since the blood-brain barrier effectively blocks reuptake), dividing the brain and its periphery into two separate compartments. Simultaneous microdialysis and blood sampling *in vivo* has provided evidence that there is sometimes a clear dissociation between release of peptides into these two compartments, and these seem to be both stimulus-dependent and peptide-specific [16]. For example, a dissociation between dendritic and axon terminal oxytocin release is evident from the effects of alpha melanocyte-stimulating hormone ( $\alpha$ -MSH). Activation of melanocortin 4 receptor receptors on oxytocin cells by  $\alpha$ -MSH mobilizes intracellular calcium and stimulates dendritic oxytocin release, but the electrical activity of the cell is inhibited, leading to less oxytocin release into the periphery [17]. Another example of dissociated release patterns is that of vasopressin release into the periphery to counteract water-loss from the kidneys in response to increased plasma osmolality. The axon terminal release of vasopressin after a systemic hypertonic saline injection increases immediately, but dendritic release of vasopressin in the SON starts only an hour later, when peripheral release is subsiding, illustrating a separation in time between release events in the dendrites and the terminals of the same neurons [18].

Whereas the SON only contains MCNs, the PVN houses many sets of functionally distinct neurons, classified into two major groups: MCNs and parvocellular neurons. Parvocellular neurosecretory neurons send their axons to the median eminence, from where they release hypophysiotropic hormones that control the function of the anterior pituitary and the major hypothalamo-pituitary axes. Parvocellular preautonomic neurons send long descending projections to sympathetic and parasympathetic centres in the brainstem and spinal cord, modulating sympathetic and parasympathetic outflows to a variety of target organs, including the heart, the peripheral vasculature and the kidneys [19-21]. In addition to neurosecretory and autonomic targets, the PVNs also include neurons that project to hierarchically higher centres in the brain, including the central amygdala, projections recently shown to modulate fear-conditioned responses [22]. These distinctive anatomical and physiological features make the PVN an ideal model to study the role of neuropeptides as signaling molecules in mediating communication within and between different neuron populations in the brain [8, 9].

## Dendritic peptide release

Modulation of neuronal function by dendritic transmitter release is a widespread phenomenon, and is specific neither to a localized part of the brain nor to a particular subtype of signalling molecule [23-26]. As mentioned above, the best-characterised sites of dendritic peptide release are the hypothalamic SON and PVN, where the MCNs release vasopressin and oxytocin from their somato-dendritic compartment. At the ultrastructural level, large dense-cored vesicles (LDCVs) are broadly distributed throughout vasopressin and oxytocin neurons and it has been shown that their contents can be released from any part of the neurons, including the cell body and especially the dendrites (Fig. 1). The first unequivocal evidence of LDCV release from dendrites came from the visualization of exocytotic profiles in electron-microscopic studies on

sympathetic and hypothalamic neurons [27-29]. Pow and Morris [29] revealed the classical LDCV morphology in the dendrites and soma of MCNs and omega-shaped fusion profiles at the plasma membrane. The authors also visualized dendritic exocytosis from oxytocin and vasopressin neurons when they treated hypothalamic tissue with tannic acid to “freeze” aggregations of the exocytosed peptide granules [29-31]. Later microsampling techniques *in vivo* confirmed and amplified the data on dendritic vasopressin and oxytocin release and revealed many aspects of its control [32].

The LDCVs often contain more than one neuropeptide, and in fact many neurons release a mixture of neuropeptides [33, 34]. For instance, vasopressin co-exists with dynorphin [35], galanin [36], pituitary adenylate cyclase activating polypeptide (PACAP) [37] and secretin in MCNs of the SON and PVN. On the other hand, oxytocin in the SON co-exists with enkephalin and dynorphin [38, 39]. Other peptides, for example apelin is also synthesized in MCNs, but it is packed and released from separate LDCVs [40].

## Mechanisms of release

### *Actin cytoskeleton*

Since peptide release from MCNs is not restricted to any particular part of the plasma membrane [29, 30], regulation of exocytosis may rely on controlling the access of the vesicles to the plasma membrane [41]. This led to the suggestion that this control may be exerted by cytoskeletal elements, as in classical endocrine cells. In addition to a network throughout the cytoplasm, the cell bodies of MCNs possess a network of filamentous protein (F-actin) beneath the plasma membrane, usually referred to as cortical F-actin. In endocrine cells, this F-actin engulfs secretory vesicles, segregating them from the plasma membrane. As F-actin undergoes fast, transient and reversible depolymerization during hormone secretion, and as areas of exocytosis have been found to be lacking F-actin, cortical F-actin has long been proposed to act as a *barrier*, restricting the movement of secretory vesicles to their release sites at the plasma membrane [42, 43].

MCNs possess F-actin structures in the subcortical regions of somata and dendrites [44, 45]. The F-actin of the somata/dendrites is rapidly and reversibly depolymerized by factors that stimulate secretion. Moreover, depolymerization of F-actin stimulates oxytocin and vasopressin release from the dendrites and acute exposure to drugs that polymerize F-actin inhibits stimulated dendritic peptide release. Thus the evoked release from the dendrites requires depolymerization of F-actin [45].

However, there is evidence that the F-actin cortex, classically viewed as a barrier that hinders the movements of LDCVs to the plasma membrane, might also play a positive role either by providing ‘tracks’ that permit docking at appropriate sites, or by spatially constraining components of the release machinery. This suggests that activation of secretion does not simply trigger the disassembly of the barrier, but rather a reorganization of F-actin, which allows the LDCVs access to the release sites and provides the structural support necessary for exocytosis [43]. In MCNs, it appears that F-actin remodelling plays a part in regulating the availability of functionally mature and readily releasable vesicles in different parts of the cell and thus is involved in the differential control of release from different parts of the cell. In contrast to neuronal synapses, release of vesicles from both the somata/dendrites and axon terminals in MCNs does not appear to occur at morphologically distinct active zones [30]. Thus, actin filaments could provide transport, tethering, barriers and support structures at different times and locations [45].

### *Exocytosis proteins*

The stimulated release of both LDCVs and synaptic vesicles involves the soluble N-ethylmaleimide sensitive factor attachment receptor (SNARE) complex, which allows the membrane of the vesicle to fuse with the plasma membrane and release its cargo into the extracellular space. There is evidence for the involvement of SNARE proteins in the release of LDCVs from dendrites, with the majority of the data arising from studies of substantia nigra dopamine cells [46-48]. Data from several other brain regions, including hippocampus [49, 50], olfactory bulb [51], cerebellum [52] and neocortex [53] indicate the requirement for SNARE variants in dendritic transmitter release.

Sensitivity of somato-dendritic release to tetanus toxin which cleaves VAMP-2 (a vesicular component of the SNARE complex) was described in isolated MCNs [54], suggesting that VAMP-2 proteins similar to

those operating in synapses may regulate dendritic exocytosis of oxytocin and vasopressin. Many SNARE proteins have been identified in the terminals of the posterior pituitary [55, 56]. However, immunofluorescence studies have shown a surprising lack of some of the core proteins, such as VAMP-2 and SNAP-25 in the somata and dendrites of the SON. Perhaps there are more members or isoforms of the existing members to be identified, but, at present, the somato-dendritic peptide release from MCNs appears to occur in the absence of the full complement of exocytosis machinery that is generally considered to be mandatory for regulated exocytosis [57].

### *Action potentials*

Exocytotic release of vasopressin and oxytocin from the axonal terminals in the posterior pituitary gland is linked to electrical activity, resulting from  $\text{Ca}^{2+}$  entry through voltage-gated channels following depolarization of the terminals by invading action potentials [58]. The available stores of small electron-lucent vesicles (ELVs) at synapses are replenished by endocytotic recycling and they are quickly re-filled with neurotransmitter by transporter-mediated uptake [59]. However, neuropeptides, which are not recycled after release have to be synthesized and the LDCVs loaded in the cell body. Compared to ELVs, LDCVs differ by requiring sustained increases in intracellular  $\text{Ca}^{2+}$  to release their contents. As a consequence, LDCVs have longer latencies to release and require stronger stimulation for exocytosis, such as, for example, bursts of electrical activity. The LDCVs also differ from ELVs in that the associated  $\text{Ca}^{2+}$ -sensor that triggers release has a higher affinity for calcium. Consequently it is not necessary for LDCVs to be located in close proximity to membrane calcium channels to undergo exocytosis, and synaptic specializations are not a prerequisite for release [60-64].

As it is the case in many neurons, the membrane properties of the dendrites support action potentials allowing them to propagate into the dendrites [65]. A rise in dendritic free  $\text{Ca}^{2+}$  content initiated by action potential back-propagation has been suggested to trigger dendritic dopamine release within the substantia nigra [46]. While action potentials may propagate into the dendrites of MCNs [66] dendritic release of vasopressin and oxytocin can occur independently of action potential firing [67, 68].

### *Role of calcium and its sources*

Calcium-dependent exocytosis represents a universal mechanism underlying release of neurotransmitters from presynaptic terminals and release of neurohormones from neuroendocrine cells. Similar to the calcium-dependent release of neuropeptides from MCNs axonal terminals in the neurohypophysis [58], dendritic release of these same neuropeptides has also been shown to be dependent on a rise in intracellular free  $\text{Ca}^{2+}$  in the dendrites [54, 69, 70].

The spatio-temporal properties and dynamics of the intracellular  $\text{Ca}^{2+}$  signal are key determinants of transmitter release in classical synapses [71]. These are in part determined by the source of  $\text{Ca}^{2+}$  and its proximity to the release machinery, as well as the different  $\text{Ca}^{2+}$  buffering mechanisms available to influence the magnitude and time course of the calcium signal. In this sense, a variety of different sources of  $\text{Ca}^{2+}$  have been shown to efficiently trigger dendritic release of oxytocin and vasopressin from MCNs.

### *Calcium channels*

A major route of entry of  $\text{Ca}^{2+}$  involved in dendritic neuropeptide release is through voltage-operated  $\text{Ca}^{2+}$  channels (VOCCs) [58, 72]. MCNs express several types of VOCCs [73], but the N-type channels appear to be particularly important for dendritic release. Although the current carried by N-type channels is comparatively small in the somata of MCNs compared to the other VOCC types or indeed the whole-cell  $\text{Ca}^{2+}$  current [74, 75], release of oxytocin from SONs is most sensitive to blockade of N-type channels. As stated above, these channels can be activated in both in somatodendritic and axonal compartments as a consequence of membrane depolarization evoked by anterograde or back-propagated action potentials [58]. However, some chemical signals, notably oxytocin and vasopressin, can themselves trigger dendritic peptide release without increasing the electrical activity of the neurons. Oxytocin- and vasopressin-neurons express oxytocin- and vasopressin-receptors, respectively [76], and the peptides act at these receptors to produce a cell-type-specific rise in intracellular  $\text{Ca}^{2+}$  concentration. For example, the response induced by vasopressin in vasopressin cells requires an influx of external  $\text{Ca}^{2+}$  through voltage-gated calcium channels, particularly of the

L-, N- and T-types [77]. The requirement of somato-dendritic release for  $\text{Ca}^{2+}$  entry through mainly L- and N-type channels has been shown for other transmitters, including dynorphin [78], dopamine [79, 80], serotonin [25] and pituitary adenylate cyclase activating polypeptide (PACAP) [70].

### NMDA receptors

Another major source of free calcium in neurons are the  $\text{Ca}^{2+}$ -permeable glutamate N-methyl-D-aspartate (NMDA) receptors. NMDA receptors are particularly important in MCNs, in which they not only influence overall MCN excitability, but also contribute to the adoption of burst-firing, optimizing in turn hormonal release from neurohypophysial terminals [81, 82]. Moreover, activation of NMDARs in MCNs results in large increases in dendritic free  $\text{Ca}^{2+}$  levels [8, 83] efficiently evoking dendritic release of both oxytocin [84] and vasopressin [8]. In addition to their conventional location at postsynaptic sites, functional NMDARs, with unique molecular and functional properties, have been also recognized to be located at extrasynaptic sites [85, 86]. In a series of recent studies, we showed the presence in MCNs of functional extrasynaptic NMDA receptors, which play a major role in regulating MCN excitability [87, 88]. Extrasynaptic NMDA receptors also contribute to increases in intracellular  $\text{Ca}^{2+}$ , and unlike synaptic NMDARs, they are selectively coupled to other  $\text{Ca}^{2+}$ -dependent signalling mechanisms, including voltage-gated potassium channels and gamma-aminobutyric acid ( $\text{GABA}_A$ ) receptors [88-90]. However, whether synaptic and extrasynaptic NMDARs selectively or differentially affect dendritic release of neuropeptides is at present unknown.

### Intracellular calcium stores

Another important source of  $\text{Ca}^{2+}$  shown to evoke and regulate dendritic release of neuropeptides are intracellular calcium stores. This is particularly the case for oxytocin autocrine effects. Binding of oxytocin to its receptors on oxytocin neurons mobilizes  $\text{Ca}^{2+}$  from intracellular stores in the endoplasmic reticulum [91]. This increase in intracellular  $\text{Ca}^{2+}$  is sufficient to induce oxytocin release from dendrites, without affecting the firing activity of neurons and without inducing release from nerve terminals [67]. Once triggered, dendritic peptide release can be self-sustaining and hence long-lasting [67]. Other agents that mobilize intracellular calcium stores, such as thapsigargin, can also evoke dendritic release of neuropeptides [67, 68, 92].

### Calcium buffering mechanisms

Intracellular  $\text{Ca}^{2+}$ -buffering mechanisms constitute additional critical factors influencing the shape and time course of intracellular  $\text{Ca}^{2+}$  signals. MCNs are endowed with numerous calcium buffering/clearance mechanisms, including plasmalemmal and endoplasmic reticulum calcium transport ATPases, the mitochondrial calcium selective uniporter (10), and  $\text{Ca}^{2+}$ -binding proteins, including calbindin and calretinin [93, 94]. Most of these mechanisms have been shown to efficiently restrain calcium transients in MCNs [83, 93, 95, 96]. Moreover, blockade of these  $\text{Ca}^{2+}$ -buffering mechanisms prolonged  $\text{K}^+$ -evoked increases in intracellular free  $\text{Ca}^{2+}$ , concomitantly enhancing somatodendritic vasopressin release [95]. Interestingly, the portfolio of available  $\text{Ca}^{2+}$ -homeostatic systems differ in somatodendritic and axonal compartments of MCNs [93, 95], further supporting the notion of independent regulation of these two neuronal compartments during neuropeptide release by MCNs.

### Calcium-dependent priming of dendritic release

In addition to directly activating dendritic release, elevation of intracellular free  $\text{Ca}^{2+}$  concentrations has another important consequence: it can *prime* dendritic stores of peptides to make them available for subsequent activity-dependent release [67]. Spike activity in oxytocin or vasopressin neurons *in vivo* does not result in measurable dendritic peptide release, but agents that mobilize  $\text{Ca}^{2+}$  from intracellular stores, such as thapsigargin or cyclopiazonic acid, or some peptides, including oxytocin itself and  $\alpha$ -MSH, consistently induce dendritic release directly [17, 67]. It seems possible that any signal that mobilizes  $\text{Ca}^{2+}$  from intracellular stores might prime dendritic secretion. Moreover, after exposure to agents that mobilize intracellular calcium, peptide release in response to many stimuli (such as osmotic stimulation, depolarization with high  $\text{K}^+$  or electrical stimulation) is dramatically potentiated. *In vitro*, this priming persists for at least 90 min. Priming involves *preparing* a system for some anticipated trigger that will come at some uncertain time in the future; it

involves making the secretory pool of the target cell available for rapid release in response to that future trigger. The mechanisms of priming in MCNs involve recruitment of vesicles from a reserve pool into a readily-releasable pool [92], probably through changes in the actin skeleton. Priming also involves recruitment of VOCCs, suggesting that a stimulus that produces an increased secretory responsiveness with an intermediate time scale (30-90min) may cause MCNs to recruit N-type calcium channels to the plasma membrane, allowing them to respond to a subsequent depolarization with a larger secretory response [75]. However, priming does not appear to require either *de novo* gene transcription or translation [97].

## **Actions of dendritically released neuropeptides**

### ***Autocrine effects***

The physiological functions of dendritically released neurotransmitters include a local autocrine effect on the neurons from which they are released, as well as effects on surrounding neurons and glia. The overall consequences can be a dramatic change in firing rate, because these autocrine effects can change both the inputs to oxytocin cells and also the way that the oxytocin cells respond to those inputs. A striking example of this is the way that dendritically released oxytocin promotes the milk ejection reflex as described below.

A far more common autocrine effect of dendritic release is *auto-inhibition*. Vasopressin neurons discharge in a characteristic phasic pattern that optimizes the efficiency of stimulus-secretion coupling at the nerve terminals. Vasopressin released from dendrites modulates this phasic activity by a predominantly inhibitory action. Interestingly, vasopressin, like oxytocin, can facilitate its own dendritic release [98]. This may explain the time dissociation between peripheral and intra-SON release of vasopressin after a hyperosmotic stimulus. Although systemic secretion of vasopressin occurs rapidly after an osmotic stimulus, the dendritic release of vasopressin evolves as a delayed and prolonged response [18]. Mimicking dendritic release by retrodialysis of vasopressin onto vasopressin neurons inhibits the vasopressin neurons by reducing their firing rate [99]. Thus, dendritic vasopressin release may activate adjacent dendrites to facilitate its own release until the local concentration has reached a threshold sufficient to hyperpolarize the neuron and/or modulate inhibitory inputs. The auto-inhibitory action of dendritic vasopressin may therefore limit the extent of systemic vasopressin secretion in response to osmotic stimuli or volume depletion.

### ***Local paracrine effects***

Exogenously applied or endogenously released oxytocin also acts on afferent nerve endings. As presynaptic oxytocin receptors are not found in the SON, this paracrine action was likely to be indirect and indeed has been shown to be mediated by oxytocin-dependent endocannabinoid release from the oxytocin neuron [100, 101]. Cannabinoid receptors (CB1) have been localized by immunohistochemistry to both excitatory and inhibitory axon terminals innervating dendrites in the SON, and the cannabinoid agonist presynaptically inhibits spontaneous excitatory and inhibitory postsynaptic currents in SON neurons recorded in slices. Thus, dendritic oxytocin release may act on oxytocin receptors leading to Ca<sup>2+</sup> release from intracellular stores and the 'on-demand' generation of endocannabinoids. The endocannabinoids pass through the membrane, diffuse and bind to presynaptic CB1 receptors, inhibiting both GABAergic and glutamatergic afferents onto MCNs. This signalling probably has a very short radius of action due to the lipophilic nature of cannabinoids. However, both oxytocin and vasopressin can spread over larger areas, effectively broadcasting their message throughout the nucleus.

An example of such longer-radius paracrine action of dendritically-released neuropeptides is highlighted by our recent study showing that dendritically-released vasopressin is able to modulate the activity of neighbouring presympathetic neurons within the PVN [8]. We found that activity-dependent dendritic release of vasopressin from MCNs resulted in a concomitant increase in the firing activity of RVLM-projecting PVN neurons. This interpopulation crosstalk involved the diffusion of vasopressin in the extracellular space, and binding and activation of V1a receptors in presympathetic neurons. We found that in contrast to conventional synaptic transmission, the efficiency and strength of this diffuse paracrine action of vasopressin was dependent on the overall extracellular levels of vasopressin (dependent in part on the average activity of the entire vasopressin population and on factors regulating vasopressin half-life in the

extracellular space) as well as the ability of vasopressin to diffuse and reach relatively distant targets (e.g., tortuosity of the extracellular space).

### *Hormone-like signals in the brain*

Oxytocin- and vasopressin-induced effects on behaviours are exerted at sites that, in some cases, richly express receptors but are innervated by few peptide-containing projections. Could dendritically released peptides act at distant brain targets to evoke long-lasting behavioural effects? Although extracellular neuropeptide concentrations differ from site to site, similar changes are often seen at widely separated sites [16]. Peptide release within the brain is not specifically targeted to synapses, and as the half-lives of peptides in the central nervous system can be up to 20 min [102], there is time for considerable movement of peptides by diffusion and bulk flow in the extracellular fluid and cerebrospinal fluid. The dendrites of MCNs project towards the brain surface and make close contact with ependymal cells that line the ventricular spaces. The reason for this may be twofold. The dendrites can register the neurochemical composition of the CSF and they can send their messages into the CSF circulation. Neuropeptides administered intracerebroventricularly lead to coherent and purposeful behaviours.

## **Physiological functions**

### *The milk ejection reflex*

Priming appears to be the key phenomenon underlying the intermittent burst discharge that oxytocin cells display in response to suckling during the milk-ejection reflex. Under basal conditions, oxytocin neurons are continuously active, but, in the pregnant animal during parturition and in the lactating animal in response to suckling, oxytocin cells discharge approximately synchronously with brief, intense bursts of action potentials; these bursts release into the circulation large boluses of oxytocin which result in intense contractions of the pregnant uterus or milk let-down from the mammary glands. For oxytocin neurons, dendritic release of oxytocin, which is up-regulated during parturition and in lactation, has an essential role in the generation of these intermittent synchronized bursts [103]. The bursting activity can be blocked by administration of oxytocin antagonists into the SON, and can be facilitated by local administration of oxytocin agonists [104].

After a priming signal, activity-dependent oxytocin release from dendrites might lead to positive-feedback coupling between oxytocin cells, producing the intense synchronized bursts observed during parturition and suckling. In each of these cases, the actions of the dendritically-released oxytocin are not restricted to the cell of origin, but are also exerted on the dendrites of other oxytocin cells, possibly to facilitate homotypic interactions.

### *Generation of multimodal neurohumoral homeostatic responses*

Control of body homeostasis by the PVN requires the generation of complex but orchestrated neurohumoral responses, generally consisting of a “neuronal” component (i.e., changes in sympathetic/parasympathetic outflows to different target organs) along with a “humoral” response, represented by the release of different neurohormones, including vasopressin, angiotensin and endothelins among others [105-107]. These neurohumoral responses generated by the PVN are critically important for the maintenance of cardiovascular and fluid balance homeostasis. A characteristic example of such an integrative homeostatic response is that following a central osmotic challenge, which evokes a coordinated increase in renal sympathetic nerve activity together with a concomitant increase in circulating levels of vasopressin. These responses are coordinated by the PVN, and result in proper adjustments in water and Na<sup>+</sup> reabsorption by the kidneys, leading in turn to reestablishment of fluid/electrolyte balance in response to the osmotic challenge [108]. We recently demonstrated that dendritically released vasopressin plays a pivotal role in this homeostatic response. We found that a central osmotic challenge evoked an increase in dendritic release of vasopressin from MCNs, which on diffusion in the extracellular space, participated in the recruitment of



neighbouring presympathetic PVN neurons. This interpopulation crosstalk resulted in turn in a proper renal sympathoexcitatory homeostatic response. Thus, dendritic release of vasopressin is a critical signalling modality contributing to the ability of the PVN to orchestrate the activity of distinct populations of neurons, and thus, the generation of multimodal homeostatic response.

### *Formation of short-term social odour memories in the olfactory system*

Both peptides evoke specific effects on behaviour [109-111]. For example, oxytocin is involved in social behaviours, including bonding and maternal behaviour, and vasopressin acts in the brain to affect social recognition and aggression. We recently identified populations of vasopressin-expressing neurons in the main and accessory olfactory bulb and in the anterior olfactory nucleus, a region of olfactory cortex that transmits and processes information in the main olfactory system [112-114]. Both vasopressin and oxytocin modulate conspecific social recognition at the level of the olfactory system and we proposed a model by which the somato-dendritic priming and release of vasopressin in main olfactory regions may facilitate the formation of short-term social odour memories [112].

#### **Acknowledgments**

We thank Dr Rafael Pineda (Edinburgh) for help with the production of the immuno-fluorescence pictures.

#### **Funding Statement**

Work was supported by grants from Biotechnology and Biological Sciences Research Council, UK (ML) and National Heart, Lung, and Blood Institute R01 HL-090948 and HL112225 (JES)

#### **Competing Interests**

We have no competing interests.

#### **Authors' Contributions**

Both authors contributed to the writing of the review.

## **Figures**

### **Figure 1) Vasopressin and oxytocin system of the hypothalamus:**

**A)** Coronal section through the rat hypothalamus at the level of the supraoptic (SON) and paraventricular nuclei (PVN); vasopressin cells are immunostained with fluorescent green and oxytocin cells with fluorescent red. **Ai)** In the SON the dendrites project towards the ventral surface of the brain where they form a dense plexus (arrow). **B)** LDCVs in a coronal section of a SON dendrite. **C)** An 'omega' fusion profile at the plasma membrane (arrow) indicates exocytosis. **D)** Close anatomical relationships among the dendrites of MCNs vasopressin (green) and retrogradely labeled presympathetic neurons from the rostral ventrolateral medulla (red) in the PVN. **Di-Dii)** Progressively higher magnification of D showing thick and varicose immunoreactive dendrites from MCN vasopressin cell dendrites in close apposition with the somata and dendrites of presympathetic neurons. Modified from [8, 16].

## References

- [1] Abbott, L.F. & Regehr, W.G. 2004 Synaptic computation. *Nature* **431**, 796-803. (doi:10.1038/nature03010).
- [2] Lisman, J.E., Raghavachari, S. & Tsien, R.W. 2007 The sequence of events that underlie quantal transmission at central glutamatergic synapses. *Nat. Rev. Neurosci.* **8**, 597-609. (doi:10.1038/nrn2191).
- [3] Mody, I., De Koninck, Y., Otis, T.S. & Soltesz, I. 1994 Bridging the cleft at GABA synapses in the brain. *Trends Neurosci.* **17**, 517-525.
- [4] Rothstein, J.D., Dykes-Hoberg, M., Pardo, C.A., Bristol, L.A., Jin, L., Kuncl, R.W., Kanai, Y., Hediger, M.A., Wang, Y., Schielke, J.P., et al. 1996 Knockout of glutamate transporters reveals a major role for astroglial transport in excitotoxicity and clearance of glutamate. *Neuron* **16**, 675-686. (doi:10.1016/S0896-6273(00)80086-0).
- [5] Zhang, Z.W., Kang, J.I. & Vaucher, E. 2011 Axonal varicosity density as an index of local neuronal interactions. *PloS One* **6**, e22543. (doi:10.1371/journal.pone.0022543).
- [6] Dawson, T.M. & Snyder, S.H. 1994 Gases as biological messengers: nitric oxide and carbon monoxide in the brain. *J. Neurosci.* **14**, 5147-5159.
- [7] Leng, G. & Ludwig, M. 2008 Neurotransmitters and peptides: whispered secrets and public announcements. *J. Physiol.* **586**, 5625-5632. (doi:10.1113/jphysiol.2008.159103).
- [8] Son, S.J., Filosa, J.A., Potapenko, E.S., Biancardi, V.C., Zheng, H., Patel, K.P., Tobin, V.A., Ludwig, M. & Stern, J.E. 2013 Dendritic peptide release mediates interpopulation crosstalk between neurosecretory and preautonomic networks. *Neuron* **78**, 1036-1049. (doi:10.1016/j.neuron.2013.04.025).
- [9] Stern, J.E. 2015 Neuroendocrine-autonomic integration in the PVN: Novel roles for dendritically released neuropeptides. *J. Neuroendocrinol.* (doi: 10.1111/jne.12252. [Epub ahead of print])
- [10] Stern, J.E. & Armstrong, W.E. 1998 Reorganization of the dendritic trees of oxytocin and vasopressin neurons of the rat supraoptic nucleus during lactation. *J. Neurosci.* **18**, 841-853.
- [11] Miyata, S. & Hatton, G.I. 2002 Activity-related, dynamic neuron-glia interactions in the hypothalamo-neurohypophysial system. *Micros. Res. Tech.* **56**, 143-157. (doi:10.1002/jemt.10012).
- [12] Hatton, G.I. 1990 Emerging concepts of structure-function dynamics in adult brain: the hypothalamo-neurohypophysial system. *Prog. Neurobiol.* **34**, 437-504.
- [13] Thome, C., Kelly, T., Yanez, A., Schultz, C., Engelhardt, M., Cambridge, S.B., Both, M., Draguhn, A., Beck, H. & Egorov, A.V. 2014 Axon-carrying dendrites convey privileged synaptic input in hippocampal neurons. *Neuron* **83**, 1418-1430. (doi:10.1016/j.neuron.2014.08.013).
- [14] Moos, F., Poulain, D.A., Rodriguez, F., Guerne, Y., Vincent, J.D. & Richard, P. 1989 Release of oxytocin within the supraoptic nucleus during the milk ejection reflex in rats. *Exp. Brain Res.* **76**, 593-602.
- [15] Wotjak, C.T., Landgraf, R. & Engelmann, M. 2008 Listening to neuropeptides by microdialysis: echoes and new sounds? *Pharmacol. Biochem. Behav.* **90**, 125-134. (doi:10.1016/j.pbb.2008.03.017).
- [16] Ludwig, M. & Leng, G. 2006 Dendritic peptide release and peptide-dependent behaviours. *Nat. Rev. Neurosci.* **7**, 126-136. (doi:10.1038/nrn1845).
- [17] Sabatier, N., Caqueneau, C., Dayanithi, G., Bull, P., Douglas, A.J., Guan, X.M., Jiang, M., Van der Ploeg, L. & Leng, G. 2003 Alpha-melanocyte-stimulating hormone stimulates oxytocin release from the dendrites of hypothalamic neurons while inhibiting oxytocin release from their terminals in the neurohypophysis. *J. Neurosci.* **23**, 10351-10358.
- [18] Ludwig, M., Callahan, M.F., Neumann, I., Landgraf, R. & Morris, M. 1994 Systemic osmotic stimulation increases vasopressin and oxytocin release within the supraoptic nucleus. *J. Neuroendocrinol.* **6**, 369-373.
- [19] Chen, Q.H. & Toney, G.M. 2003 Identification and characterization of two functionally distinct groups of spinal cord-projecting paraventricular nucleus neurons with sympathetic-related activity. *Neuroscience* **118**, 797-807.
- [20] Stocker, S.D., Keith, K.J. & Toney, G.M. 2004 Acute inhibition of the hypothalamic paraventricular nucleus decreases renal sympathetic nerve activity and arterial blood pressure in water-deprived rats. *Am. J. Physiol.* **286**, R719-725. (doi:10.1152/ajpregu.00494.2003).
- [21] Zhang, K. & Patel, K.P. 1998 Effect of nitric oxide within the paraventricular nucleus on renal sympathetic nerve discharge: role of GABA. *Am. J. Physiol.* **275**, R728-734.
- [22] Knobloch, H.S., Charlet, A., Hoffmann, L.C., Eliava, M., Khrulev, S., Cetin, A.H., Osten, P., Schwarz, M.K., Seeburg, P.H., Stoop, R., et al. 2012 Evoked axonal oxytocin release in the central amygdala attenuates fear response. *Neuron* **73**, 553-566. (doi:10.1016/j.neuron.2011.11.030).
- [23] Ludwig, M. & Pittman, Q.J. 2003 Talking back: dendritic neurotransmitter release. *Trends Neurosci.* **26**, 255-261. (doi:10.1016/S0166-2236(03)00072-9).
- [24] Kennedy, M.J. & Ehlers, M.D. 2011 Mechanisms and function of dendritic exocytosis. *Neuron* **69**, 856-875. (doi:10.1016/j.neuron.2011.02.032).
- [25] Trueta, C. & De-Miguel, F.F. 2012 Extrasynaptic exocytosis and its mechanisms: a source of molecules mediating volume transmission in the nervous system. *Front. Physiol.* **3**, 319. (doi:10.3389/fphys.2012.00319).
- [26] Cox, C.L. 2014 Complex regulation of dendritic transmitter release from thalamic interneurons. *Curr. Opin. Neurobiol.* **29C**, 126-132. (doi:10.1016/j.conb.2014.07.004).
- [27] Zaidi, Z.F. & Matthews, M.R. 1997 Exocytotic release from neuronal cell bodies, dendrites and nerve terminals in sympathetic ganglia of the rat, and its differential regulation. *Neuroscience* **80**, 861-891.
- [28] Zaidi, Z.F. & Matthews, M.R. 1999 Stimulant-induced exocytosis from neuronal somata, dendrites, and newly formed synaptic nerve terminals in chronically decentralized sympathetic ganglia of the rat. *J. Comp. Neurol.* **415**, 121-143.
- [29] Pow, D.V. & Morris, J.F. 1989 Dendrites of hypothalamic magnocellular neurons release neurohypophysial peptides by exocytosis. *Neuroscience* **32**, 435-439.
- [30] Morris, J.F. & Pow, D.V. 1991 Widespread release of peptides in the central nervous system: quantitation of tannic acid-captured exocytoses. *Anat. Rec.* **231**, 437-445. (doi:10.1002/ar.1092310406).

- [31] Morris, J.F. & Ludwig, M. 2004 Magnocellular dendrites: prototypic receiver/transmitters. *J. Neuroendocrinol.* **16**, 403-408. (doi:10.1111/j.0953-8194.2004.01182.x).
- [32] Ludwig, M. 1998 Dendritic release of vasopressin and oxytocin. *J. Neuroendocrinol.* **10**, 881-895.
- [33] Merighi, A. 2002 Costorage and coexistence of neuropeptides in the mammalian CNS. *Prog. Neurobiol.* **66**, 161-190.
- [34] Meister, B., Villar, M.J., Ceccatelli, S. & Hokfelt, T. 1990 Localization of chemical messengers in magnocellular neurons of the hypothalamic supraoptic and paraventricular nuclei: an immunohistochemical study using experimental manipulations. *Neuroscience* **37**, 603-633.
- [35] Watson, S.J., Akil, H., Fischli, W., Goldstein, A., Zimmerman, E., Nilaver, G. & van wimersma Griedanus, T.B. 1982 Dynorphin and vasopressin: common localization in magnocellular neurons. *Science* **216**, 85-87.
- [36] Landry, M., Vila-Porcile, E., Hokfelt, T. & Calas, A. 2003 Differential routing of coexisting neuropeptides in vasopressin neurons. *Eur. J. Neurosci.* **17**, 579-589.
- [37] Gillard, E.R., Leon-Olea, M., Mucio-Ramirez, S., Coburn, C.G., Sanchez-Islas, E., de Leon, A., Mussenden, H., Bause, L.G., Pittman, Q.J. & Curras-Collazo, M.C. 2006 A novel role for endogenous pituitary adenylate cyclase activating polypeptide in the magnocellular neuroendocrine system. *Endocrinology* **147**, 791-803. (doi:10.1210/en.2005-1103).
- [38] Martin, R., Moll, U. & Voigt, K.H. 1983 An attempt to characterize by immunocytochemical methods the enkephalin-like material in oxytocin endings of the rat neurohypophysis. *Life Sci.* **33 Suppl 1**, 69-72.
- [39] Eriksson, M., Ceccatelli, S., Uvnas-Moberg, K., Iadarola, M. & Hokfelt, T. 1996 Expression of Fos-related antigens, oxytocin, dynorphin and galanin in the paraventricular and supraoptic nuclei of lactating rats. *Neuroendocrinology* **63**, 356-367.
- [40] De Mota, N., Reaux-Le Goazigo, A., El Messari, S., Chartrel, N., Roesch, D., Dujardin, C., Kordon, C., Vaudry, H., Moos, F. & Llorens-Cortes, C. 2004 Apelin, a potent diuretic neuropeptide counteracting vasopressin actions through inhibition of vasopressin neuron activity and vasopressin release. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 10464-10469. (doi:10.1073/pnas.0403518101).
- [41] Morgan, A. 1995 Exocytosis. *Essays Biochem.* **30**, 77-95.
- [42] Vitale, M.L., Seward, E.P. & Trifaro, J.M. 1995 Chromaffin cell cortical actin network dynamics control the size of the release-ready vesicle pool and the initial rate of exocytosis. *Neuron* **14**, 353-363. (doi:0896-6273(95)90291-0 [pii]).
- [43] Dillon, C. & Goda, Y. 2005 The actin cytoskeleton: integrating form and function at the synapse. *Annu. Rev. Neurosci.* **28**, 25-55. (doi:10.1146/annurev.neuro.28.061604.135757).
- [44] Wang, Y.F. & Hatton, G.I. 2006 Mechanisms underlying oxytocin-induced excitation of supraoptic neurons: prostaglandin mediation of actin polymerization. *J. Neurophysiol.* **95**, 3933-3947. (doi:10.1152/jn.01267.2005).
- [45] Tobin, V.A. & Ludwig, M. 2007 The role of the actin cytoskeleton in oxytocin and vasopressin release from rat supraoptic nucleus neurons. *J. Physiol.* **582**, 1337-1348. (doi:10.1113/jphysiol.2007.132639).
- [46] Bergquist, F. & Ludwig, M. 2008 Dendritic transmitter release: a comparison of two model systems. *J. Neuroendocrinol.* **20**, 677-686. (doi:10.1111/j.1365-2826.2008.01714.x).
- [47] Witkovsky, P., Patel, J.C., Lee, C.R. & Rice, M.E. 2009 Immunocytochemical identification of proteins involved in dopamine release from the somatodendritic compartment of nigral dopaminergic neurons. *Neuroscience* **164**, 488-496. (doi:10.1016/j.neuroscience.2009.08.017).
- [48] Ovsepien, S.V. & Dolly, J.O. 2011 Dendritic SNAREs add a new twist to the old neuron theory. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 19113-19120. (doi:10.1073/pnas.1017235108 [pii] 10.1073/pnas.1017235108).
- [49] Maletic-Savatic, M., Koothan, T. & Malinow, R. 1998 Calcium-evoked dendritic exocytosis in cultured hippocampal neurons. Part II: mediation by calcium/calmodulin-dependent protein kinase II. *J. Neurosci.* **18**, 6814-6821.
- [50] Maletic-Savatic, M. & Malinow, R. 1998 Calcium-evoked dendritic exocytosis in cultured hippocampal neurons. Part I: trans-Golgi network-derived organelles undergo regulated exocytosis. *J. Neurosci.* **18**, 6803-6813.
- [51] Matsutani, S. & Yamamoto, N. 2004 Postnatal development of dendritic spines on olfactory bulb granule cells in rats. *J. Comp. Neurol.* **473**, 553-561. (doi:10.1002/cne.20107).
- [52] Duguid, I.C., Pankratov, Y., Moss, G.W. & Smart, T.G. 2007 Somatodendritic release of glutamate regulates synaptic inhibition in cerebellar Purkinje cells via autocrine mGluR1 activation. *J. Neurosci.* **27**, 12464-12474. (doi:10.1523/JNEUROSCI.0178-07.2007).
- [53] Zilberter, Y., Harkany, T. & Holmgren, C.D. 2005 Dendritic release of retrograde messengers controls synaptic transmission in local neocortical networks. *Neuroscientist* **11**, 334-344. (doi:10.1177/1073858405275827).
- [54] de Kock, C.P., Wierda, K.D., Bosman, L.W., Min, R., Koksmas, J.J., Mansvelder, H.D., Verhage, M. & Brussaard, A.B. 2003 Somatodendritic secretion in oxytocin neurons is upregulated during the female reproductive cycle. *J. Neurosci.* **23**, 2726-2734. (doi:10.1523/JNEUROSCI.2726-03.2003).
- [55] Jurgutis, P., Shuang, R., Fletcher, A. & Stuenkel, E.L. 1996 Characterization and distribution of SNARE proteins at neuroendocrine nerve endings. *Neuroendocrinology* **64**, 379-392.
- [56] Zhang, Z., Bhalla, A., Dean, C., Chapman, E.R. & Jackson, M.B. 2009 Synaptotagmin IV: a multifunctional regulator of peptidergic nerve terminals. *Nat. Neurosci.* **12**, 163-171. (doi:10.1038/nn.2252).
- [57] Tobin, V., Schwab, Y., Lelos, N., Onaka, T., Pittman, Q.J. & Ludwig, M. 2012 Expression of exocytosis proteins in rat supraoptic nucleus neurons. *J. Neuroendocrinol.* **24**, 629-641. (doi:10.1111/j.1365-2826.2011.02237.x).
- [58] Fisher, T.E. & Bourque, C.W. 1996 Calcium-channel subtypes in the somata and axon terminals of magnocellular neurosecretory cells. *Trends Neurosci.* **19**, 440-444. (doi:10.1016/j.tins.1996.05.004 [pii]).
- [59] Prior, I.A. & Clague, M.J. 1997 Glutamate uptake occurs at an early stage of synaptic vesicle recycling. *Curr. Biol.* **7**, 353-356.
- [60] An, S. & Zenisek, D. 2004 Regulation of exocytosis in neurons and neuroendocrine cells. *Curr. Opin. Neurobiol.* **14**, 522-530. (doi:10.1016/j.conb.2004.08.008).
- [61] Baraban, S.C. & Tallent, M.K. 2004 Interneuron Diversity series: Interneuronal neuropeptides--endogenous regulators of neuronal excitability. *Trends Neurosci.* **27**, 135-142. (doi:10.1016/j.tins.2004.01.008).
- [62] Mansvelder, H.D. & Kits, K.S. 2000 Calcium channels and the release of large dense core vesicles from neuroendocrine cells: spatial organization and functional coupling. *Prog. Neurobiol.* **62**, 427-441.
- [63] Martin, T.F. 2003 Tuning exocytosis for speed: fast and slow modes. *Biochim. Biophys. Acta* **1641**, 157-165.
- [64] Salio, C., Lossi, L., Ferrini, F. & Merighi, A. 2006 Neuropeptides as synaptic transmitters. *Cell Tissue Res.* **326**, 583-598. (doi:10.1007/s00441-006-0268-3).

- [65] Stuart, G., Spruston, N., Sakmann, B. & Häusser, M. 1997 Action potential initiation and backpropagation in neurons of the mammalian CNS. *Trends Neurosci.* **20**, 125-131.
- [66] Bains, J.S. & Ferguson, A.V. 1999 Activation of N-methyl-D-aspartate receptors evokes calcium spikes in the dendrites of rat hypothalamic paraventricular nucleus neurons. *Neuroscience* **90**, 885-891.
- [67] Ludwig, M., Sabatier, N., Bull, P.M., Landgraf, R., Dayanithi, G. & Leng, G. 2002 Intracellular calcium stores regulate activity-dependent neuropeptide release from dendrites. *Nature* **418**, 85-89. (doi:10.1038/nature00822).
- [68] Ludwig, M., Bull, P.M., Tobin, V.A., Sabatier, N., Landgraf, R., Dayanithi, G. & Leng, G. 2005 Regulation of activity-dependent dendritic vasopressin release from rat supraoptic neurones. *J. Physiol.* **564**, 515-522. (doi:10.1113/jphysiol.2005.083931).
- [69] Neumann, I., Russell, J.A. & Landgraf, R. 1993 Oxytocin and vasopressin release within the supraoptic and paraventricular nuclei of pregnant, parturient and lactating rats: a microdialysis study. *Neuroscience* **53**, 65-75. (doi:0306-4522(93)90285-N [pii]).
- [70] Shibuya, I., Noguchi, J., Tanaka, K., Harayama, N., Inoue, U., Kabashima, N., Ueta, Y., Hattori, Y. & Yamashita, H. 1998 PACAP increases the cytosolic Ca<sup>2+</sup> concentration and stimulates somatodendritic vasopressin release in rat supraoptic neurons. *J. Neuroendocrinol.* **10**, 31-42.
- [71] Meinrenken, C.J., Borst, J.G. & Sakmann, B. 2003 Local routes revisited: the space and time dependence of the Ca<sup>2+</sup> signal for phasic transmitter release at the rat calyx of Held. *J. Physiol.* **547**, 665-689. (doi:10.1113/jphysiol.2002.032714).
- [72] Wang, D. & Fisher, T.E. 2014 Expression of CaV 2.2 and splice variants of CaV 2.1 in oxytocin- and vasopressin-releasing supraoptic neurones. *J. Neuroendocrinol.* **26**, 100-110. (doi:10.1111/jne.12127).
- [73] Foehring, R.C. & Armstrong, W.E. 1996 Pharmacological dissection of high-voltage-activated Ca<sup>2+</sup> current types in acutely dissociated rat supraoptic magnocellular neurons. *J. Neurophysiol.* **76**, 977-983.
- [74] Joux, N., Chevaléyre, V., Alonso, G., Boissin-Agasse, L., Moos, F.C., Desarmenien, M.G. & Hussy, N. 2001 High voltage-activated Ca<sup>2+</sup> currents in rat supraoptic neurones: biophysical properties and expression of the various channel  $\alpha$ 1 subunits. *J. Neuroendocrinol.* **13**, 638-649. (doi:jne679 [pii]).
- [75] Tobin, V.A., Douglas, A.J., Leng, G. & Ludwig, M. 2011 The involvement of voltage-operated calcium channels in somato-dendritic oxytocin release. *PLoS One* **6**, e25366. (doi:10.1371/journal.pone.0025366).
- [76] Freund-Mercier, M.J., Stoeckel, M.E. & Klein, M.J. 1994 Oxytocin receptors on oxytocin neurones: histoautoradiographic detection in the lactating rat. *J. Physiol.* **480**, 155-161.
- [77] Sabatier, N., Richard, P. & Dayanithi, G. 1997 L-, N- and T- but neither P- nor Q-type Ca<sup>2+</sup> channels control vasopressin-induced Ca<sup>2+</sup> influx in magnocellular vasopressin neurones isolated from the rat supraoptic nucleus. *J. Physiol.* **503**, 253-268.
- [78] Simmons, M.L., Terman, G.W., Gibbs, S.M. & Chavkin, C. 1995 L-type calcium channels mediate dynorphin neuropeptide release from dendrites but not axons of hippocampal granule cells. *Neuron* **14**, 1265-1272. (doi:0896-6273(95)90273-2 [pii]).
- [79] Kim, Y., Park, M.K. & Chung, S. 2009 Regulation of somatodendritic dopamine release by corticotropin-releasing factor via the inhibition of voltage-operated Ca<sup>2+</sup> channels. *Neurosci. Lett.* **465**, 31-35. (doi:S0304-3940(09)01173-2 [pii] 10.1016/j.neulet.2009.08.066).
- [80] Mendez, J.A., Bourque, M.J., Fasano, C., Kortleven, C. & Trudeau, L.E. 2011 Somatodendritic dopamine release requires synaptotagmin 4 and 7 and the participation of voltage-gated calcium channels. *J. Biol. Chem.* **286**, 23928-23937. (doi:10.1074/jbc.M111.218032).
- [81] Hu, B. & Bourque, C.W. 1992 NMDA receptor-mediated rhythmic bursting activity in rat supraoptic nucleus neurones in vitro. *J. Physiol.* **458**, 667-687.
- [82] Nissen, R., Hu, B. & Renaud, L.P. 1995 Regulation of spontaneous phasic firing of rat supraoptic vasopressin neurones in vivo by glutamate receptors. *J. Physiol.* **484**, 415-424.
- [83] Stern, J.E. & Potapenko, E.S. 2013 Enhanced NMDA receptor-mediated intracellular calcium signaling in magnocellular neurosecretory neurons in heart failure rats. *Am. J. Physiol.* **305**, R414-422. (doi:10.1152/ajpregu.00160.2013).
- [84] de Kock, C.P., Burnashev, N., Lodder, J.C., Mansvelder, H.D. & Brussaard, A.B. 2004 NMDA receptors induce somatodendritic secretion in hypothalamic neurones of lactating female rats. *J. Physiol.* **561**, 53-64. (doi:10.1113/jphysiol.2004.069005).
- [85] Sah, P., Hestrin, S. & Nicoll, R.A. 1989 Tonic activation of NMDA receptors by ambient glutamate enhances excitability of neurons. *Science* **246**, 815-818.
- [86] Le Meur, K., Galante, M., Angulo, M.C. & Audinat, E. 2007 Tonic activation of NMDA receptors by ambient glutamate of non-synaptic origin in the rat hippocampus. *J. Physiol.* **580**, 373-383. (doi:10.1113/jphysiol.2006.123570).
- [87] Fleming, T.M., Scott, V., Naskar, K., Joe, N., Brown, C.H. & Stern, J.E. 2011 State-dependent changes in astrocyte regulation of extrasynaptic NMDA receptor signalling in neurosecretory neurons. *J. Physiol.* **589**, 3929-3941. (doi:10.1113/jphysiol.2011.207340).
- [88] Naskar, K. & Stern, J.E. 2014 A functional coupling between extrasynaptic NMDA receptors and A-type K<sup>+</sup> channels under astrocyte control regulates hypothalamic neurosecretory neuronal activity. *J. Physiol.* **592**, 2813-2827. (doi:10.1113/jphysiol.2014.270793).
- [89] Potapenko, E.S., Biancardi, V.C., Florschütz, R.M., Ryu, P.D. & Stern, J.E. 2011 Inhibitory-excitatory synaptic balance is shifted toward increased excitation in magnocellular neurosecretory cells of heart failure rats. *J. Neurophysiol.* **106**, 1545-1557. (doi:10.1152/jn.00218.2011).
- [90] Potapenko, E.S., Biancardi, V.C., Zhou, Y. & Stern, J.E. 2013 Astrocytes modulate a postsynaptic NMDA-GABA<sub>A</sub>-receptor crosstalk in hypothalamic neurosecretory neurons. *J. Neurosci.* **33**, 631-640. (doi:10.1523/JNEUROSCI.3936-12.2013).
- [91] Lambert, R.C., Dayanithi, G., Moos, F.C. & Richard, P. 1994 A rise in the intracellular Ca<sup>2+</sup> concentration of isolated rat supraoptic cells in response to oxytocin. *J. Physiol.* **478**, 275-287.
- [92] Tobin, V.A., Hurst, G., Norrie, L., Dal Rio, F.P., Bull, P.M. & Ludwig, M. 2004 Thapsigargin-induced mobilization of dendritic dense-cored vesicles in rat supraoptic neurons. *Eur. J. Neurosci.* **19**, 2909-2912. (doi:10.1111/j.1460-9568.2004.03388.x).
- [93] Dayanithi, G., Forostyak, O., Ueta, Y., Verkhratsky, A. & Toescu, E.C. 2012 Segregation of calcium signalling mechanisms in magnocellular neurones and terminals. *Cell Calcium* **51**, 293-299. (doi:10.1016/j.ceca.2012.02.002).
- [94] Miyata, S., Khan, A.M. & Hatton, G.I. 1998 Colocalization of calretinin and calbindin-D28k with oxytocin and vasopressin in rat supraoptic nucleus neurons: a quantitative study. *Brain Res.* **785**, 178-182.
- [95] Komori, Y., Tanaka, M., Kuba, M., Ishii, M., Abe, M., Kitamura, N., Verkhratsky, A., Shibuya, I. & Dayanithi, G. 2010 Ca(2+) homeostasis, Ca(2+) signalling and somatodendritic vasopressin release in adult rat supraoptic nucleus neurones. *Cell Calcium* **48**, 324-332. (doi:10.1016/j.ceca.2010.10.002).

- [96] Shaw, F.D. & Morris, J.F. 1980 Calcium localization in the rat neurohypophysis. *Nature* **287**, 56-58.
- [97] Tobin, V., Leng, G. & Ludwig, M. 2012 The involvement of actin, calcium channels and exocytosis proteins in somato-dendritic oxytocin and vasopressin release. *Front. Physiol.* **3**, 261. (doi:10.3389/fphys.2012.00261).
- [98] Wotjak, C.T., Ludwig, M. & Landgraf, R. 1994 Vasopressin facilitates its own release within the rat supraoptic nucleus in vivo. *Neuroreport* **5**, 1181-1184.
- [99] Ludwig, M. & Leng, G. 1997 Autoinhibition of supraoptic nucleus vasopressin neurons in vivo: a combined retrodialysis/electrophysiological study in rats. *Eur. J. Neurosci.* **9**, 2532-2540.
- [100] Kombian, S.B., Mougnot, D. & Pittman, Q.J. 1997 Dendritically released peptides act as retrograde modulators of afferent excitation in the supraoptic nucleus in vitro. *Neuron* **19**, 903-912.
- [101] Oliet, S.H., Baimoukhametova, D.V., Piet, R. & Bains, J.S. 2007 Retrograde regulation of GABA transmission by the tonic release of oxytocin and endocannabinoids governs postsynaptic firing. *J. Neurosci.* **27**, 1325-1333. (doi:10.1523/JNEUROSCI.2676-06.2007).
- [102] Mens, W.B., Witter, A. & van Wimersma Greidanus, T.B. 1983 Penetration of neurohypophyseal hormones from plasma into cerebrospinal fluid (CSF): half-times of disappearance of these neuropeptides from CSF. *Brain Res.* **262**, 143-149.
- [103] Rossoni, E., Feng, J., Tirozzi, B., Brown, D., Leng, G. & Moos, F. 2008 Emergent synchronous bursting of oxytocin neuronal network. *PLoS Comp. Biol.* **4**, e1000123. (doi:10.1371/journal.pcbi.1000123).
- [104] Lambert, R.C., Moos, F.C. & Richard, P. 1993 Action of endogenous oxytocin within the paraventricular or supraoptic nuclei: a powerful link in the regulation of the bursting pattern of oxytocin neurons during the milk-ejection reflex in rats. *Neuroscience* **57**, 1027-1038.
- [105] Swanson, L.W. & Kuypers, H.G. 1980 The paraventricular nucleus of the hypothalamus: cytoarchitectonic subdivisions and organization of projections to the pituitary, dorsal vagal complex, and spinal cord as demonstrated by retrograde fluorescence double-labeling methods. *J. Comp. Neurol.* **194**, 555-570. (doi:10.1002/cne.901940306).
- [106] Patel, K.P. & Zhang, K. 1996 Neurohumoral activation in heart failure: role of paraventricular nucleus. *Clin. Exp. Pharmacol. Physiol.* **23**, 722-726.
- [107] Stern, J.E. & Filosa, J.A. 2013 Bidirectional neuro-glial signaling modalities in the hypothalamus: role in neurohumoral regulation. *Auton. Neurosci.* **175**, 51-60. (doi:10.1016/j.autneu.2012.12.009).
- [108] Bourque, C.W. 2008 Central mechanisms of osmosensation and systemic osmoregulation. *Nat. Rev. Neurosci.* **9**, 519-531. (doi:10.1038/nrn2400).
- [109] Neumann, I.D. & Landgraf, R. 2012 Balance of brain oxytocin and vasopressin: implications for anxiety, depression, and social behaviors. *Trends Neurosci.* **35**, 649-659. (doi:10.1016/j.tins.2012.08.004).
- [110] Insel, T.R. 2010 The challenge of translation in social neuroscience: a review of oxytocin, vasopressin, and affiliative behavior. *Neuron* **65**, 768-779. (doi:10.1016/j.neuron.2010.03.005).
- [111] Stoop, R. 2014 Neuromodulation by oxytocin and vasopressin in the central nervous system as a basis for their rapid behavioral effects. *Curr. Opin. Neurobiol.* **29**, 187-193. (doi:10.1016/j.conb.2014.09.012).
- [112] Tobin, V.A., Hashimoto, H., Wacker, D.W., Takayanagi, Y., Langnaese, K., Caquineau, C., Noack, J., Landgraf, R., Onaka, T., Leng, G., et al. 2010 An intrinsic vasopressin system in the olfactory bulb is involved in social recognition. *Nature* **464**, 413-417. (doi:10.1038/nature08826).
- [113] Wacker, D.W., Tobin, V.A., Noack, J., Bishop, V.R., Duszkievicz, A.J., Engelmann, M., Meddle, S.L. & Ludwig, M. 2010 Expression of early growth response protein 1 in vasopressin neurones of the rat anterior olfactory nucleus following social odour exposure. *J. Physiol.* **588**, 4705-4717. (doi:10.1113/jphysiol.2010.196139).
- [114] Wacker, D.W. & Ludwig, M. 2012 Vasopressin, oxytocin, and social odor recognition. *Horm. Behav.* **61**, 259-265. (doi:10.1016/j.yhbeh.2011.08.014).